


**OPTIMISATION OF OLEANOLIC ACID EXTRACTION FROM
'POKOK BUNGA TAHI AYAM' (*LANTANA CAMARA*) ROOTS
BY METHANOL-ETHYL ACETATE SOLVENT USING
RESORSE SURFACE METHODOLOGY**

RASIMAH MAHAT


**Final Year Project Report Submitted in
Partial Fulfillment of the Requirements for the
Degree of Bachelor of Sciences (Hons.) Applied Chemistry
in the Faculty of Applied Sciences
Universiti Teknologi MARA**

NOVEMBER 2008

This Final Year Report entitled “ **Optimisation of Oleanolic Acid Extraction From *Lantana camara* (Pokok Bunga Tahi Ayam) Roots by Methanol-Ethyl Acetate Solvent Using Response Surface Methodology**” was submitted by Rasimah Mahat, in partial fulfillment of the requirements for the Degree of Bachelor of Science (Hons.) Applied Science, in the Faculty of Applied Sciences, and was approved by



Assoc. Prof. Dr. Zainal Samicho
Supervisor
B. SC. (Hons.) Food Science and Technology
Faculty of Applied Sciences
Universiti Teknologi MARA
40450 Shah Alam
Selangor



Ms. Sabrina binti M. Yahya
Project Coordinator
B.Sc. (Hons.) Applied Chemistry
Faculty of Applied Sciences
Universiti Teknologi MARA
40450 Shah Alam
Selangor



Dr. Yusairie bin Mohd
Head of Programme
B. Sc. (Hons) Applied Chemistry
Faculty of Applied Sciences
Universiti Teknologi MARA
40450 Shah Alam
Selangor

Date : 28 NOV 2008

ACKNOWLEDGEMENTS

I would like to express my deep and sincere gratitude to my supervisor, Associate Professor Dr. Zainal Samicho, Lecturer Food Technology, Universiti Teknologi MARA Malaysia, Shah Alam. His wide knowledge and his logical way of thinking have been of great value for me. His understanding, encouraging and personal guidance in all the time of research and writing of this thesis have strengthen my spirit as well as broaden my knowledge. Also his wife whose willing to take care my sample.

I would like to express my gratitude to all those who gave me the possibility to complete this thesis. I want to thank to Mr. Muhd Fahmy Bin Hj. Ishak, owner of the Aneka Nursery, Kg. Sg Udang Klang, Selangor for his encouragement helped pertaining to find my sample.

I would like to wish an acknowledge to all the laboratory staff whose giving me a great cooperation by providing relevant information pertaining to the instrument handling and willing to stay back and let me used the lab over the office hour.

I would also like to thanks to my beloved parents, Mr. Mahat bin Salleh and Seti Martin bt Martin who continuously gave me the spirit, strength and also encourage me to always do my best and to not give up. Without their care, attention and love, maybe it is difficult for me to perform and do my thesis as good as what I had done.

I kindly appreciate my beloved friends and other parties for their help either directly or indirectly during my way to complete this research. Thanks to them. None of this would have been hard without their support.

Rasimah Binti Mahat

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
TABLE OF CONTENTS	iv
LIST OF TABLES	vi
LIST OF FIGURES	vii
LIST OF ABBREVIATION	viii
ABSTRACT	ix
ABSTRAK	x
 CHAPTER 1 INTRODUCTION	
1.1 Background	1
1.2 Significant of study	1
1.3 Objectives of study	2
 CHAPTER 2 LITERATURE REVIEW	
2.1 <i>Lantana camara</i>	3
2.1.1 Classification	3
2.1.2 Discription	3
2.1.3 Bioactive principle in <i>Lantana camara</i>	5
2.2 Oleanolic acid	6
2.2.1 Molecular formula	6
2.2.2 The IUPAC name of oleanolic acid	6
2.2.3 Molecular structure	6
2.2.4 Physical properties	7
2.2.5 Chemical compositions	7
2.2.6 Pharmacological properties	9
2.2.6.1 Hepatoprotective effects	10
2.2.6.2 Anti-inflammatory effects	11
2.2.6.3 Anti-tumor activity	12
2.2.6.4 Anti-hyperlipidermic effects	13
2.2.6.5 Anti-ulcers effects	13
2.3 Extraction of oleanolic acid	14
 CHAPTER 3 METHODOLOGY	
3.1 Material	15
3.1.1 Raw material	15
3.1.2 Reagents	15
3.1.3 Apparatus	16

ABSTRACT

OPTIMISATION OF OLEANOLIC ACID EXTRACTION FROM 'POKOK BUNGA TAHI AYAM' (*LANTANA CAMARA*) ROOTS BY USING RESPONSE SURFACE METHODOLOGY (RSM)

Lantana camara is an ornamental plant used in tradition medicine for the treatment of various diseases. The roots of *L. camara* is a rich of oleanolic acid which has shown anti-inflammatory, hepatoprotective, antitumor, antioxidant and anti-hyperliperemic activity. Optimisation of various extraction parameters using response surface methodology (RSM) was performed to asses maximum yield of oleanolic acid from *L.camara* roots. MINITAB software version 14 was used to identify the significant effect of various extraction parameters such as concentration of solvent-solid ratio (mL:g), extraction temperature and extraction time. Among the three variables tested, concentration of solvent-solid ratio (mL:g) and extraction temperature were found to have significant effect on oleanolic acid extraction. Optimum level of the significant variables were determined by using central composite design. The optimum condition for extraction of oleanolic acid was found at 30 mL/g of concentration of solvent-solid ratio, 60°C of extraction temperature and 30 minutes of extraction time. At optimum extraction parameters, the optimum yield of oleanolic acid obtained experimentally (1.36609 %) was found to be very close to its predicted value determined by (1.2915%) MINITAB software version 14.